Bull. Math. Soc. Sci. Math. Roumanie Tome 54(102) No. 2, 2011, 109–117

On the set of mild solutions of a nonconvex nonclosed evolution inclusion

by Aurelian Cernea

Abstract

We consider a nonconvex and nonclosed evolution inclusion and we prove the arcwise connectedness of the set of its mild solutions.

Key Words: Set-valued contraction, fixed point, mild solution. **2010 Mathematics Subject Classification**: Primary 34A60; Secondary 45D99.

1 Introduction

This paper is concerned with the following differential inclusion

$$x'(t) \in A(t)x(t) + \int_0^t K(t,s)F(s,x(s),H(s,x(s)))ds, \quad x(0) = x_0,$$
(1.1)

where $F : [0, T] \times X \times X \to \mathcal{P}(X), H : [0, T] \times X \to \mathcal{P}(X)$ are set-valued maps, X is a separable Banach space, A(t) is the infinitesimal generator of a strongly continuous evolution system of a two parameter family $\{G(t, \tau), t \ge 0, \tau \ge 0\}$ of bounded linear operators of X into X, $D = \{(t, s) \in [0, T] \times [0, T]; t \ge s\},$ $K(., .) : D \to \mathbf{R}$ is continuous and $x_0 \in X$.

When F does not depend on the last variable (1.1) reduces to

$$x'(t) \in A(t)x(t) + \int_0^t K(t,s)F(s,x(s))ds, \quad x(0) = x_0,$$
(1.2)

Existence results and qualitative properties of the mild solutions of problem (1.2) may be found in [1-7] etc. In all these papers the set-valued map F is assumed to be at least closed-valued. Such an assumption is quite natural in order to obtain good properties of the solution set, but it is interesting to investigate the problem when the right-hand side of the multivalued equation may have nonclosed values.

Following the approach in [13] we consider problem (1.1), where F and H are closed-valued multifunctions Lipschitzian with respect to the second variable and F is contractive in the third variable. Obviously, the right-hand side of the differential inclusion in (1.1) is in general neither convex nor closed. We prove the arcwise connectedness of the solution set to (1.1). The main tool is a result ([11,12]) concerning the arcwise connectedness of the fixed point set of a class of nonconvex nonclosed set-valued contractions. We note that this idea was already used for similar results concerning other classes of differential inclusions ([8-10]).

The paper is organized as follows: in Section 2 we recall some preliminary results that we use in the sequel and in Section 3 we prove our main result.

2 Preliminaries

Let Z be a metric space with the distance d_Z and let 2^Z be the family of all nonempty closed subsets of Z. For $a \in Z$ and $A, B \in 2^Z$ set $d_Z(a, B) =$ $\inf_{b \in B} d_Z(a, b)$ and $d_Z^*(A, B) = \sup_{a \in A} d_Z(a, B)$. Denote by D_Z the Pompeiu-Hausdorff generalized metric on 2^Z defined by

$$D_Z(A,B) = \max\{d_Z^*(A,B), d_Z^*(B,A)\}, A, B \in 2^Z.$$

In what follows, when the product $Z = Z_1 \times Z_2$ of metric spaces $Z_i, i = 1, 2$, is considered, it is assumed that Z is equipped with the distance $d_Z((z_1, z_2), (z'_1, z'_2)) = \sum_{i=1}^2 d_{Z_i}(z_i, z'_i)$. Let X be a nonempty set and let $F: X \to 2^Z$ be a set-valued map from X

Let X be a nonempty set and let $F: X \to 2^Z$ be a set-valued map from X to Z. The range of F is the set $F(X) = \bigcup_{x \in X} F(x)$. Let (X, \mathcal{F}) be a measurable space. The multifunction $F: X \to 2^Z$ is called measurable if $F^{-1}(\Omega) \in \mathcal{F}$ for any open set $\Omega \subset Z$, where $F^{-1}(\Omega) = \{x \in X; F(x) \cap \Omega \neq \emptyset\}$. Let (X, d_X) be a metric space. The multifunction F is called Hausdorff continuous if for any $x_0 \in X$ and every $\epsilon > 0$ there exists $\delta > 0$ such that $x \in X, d_X(x, x_0) < \delta$ implies $D_Z(F(x), F(x_0)) < \epsilon$.

Let (P, \mathcal{F}, μ) be a finite, positive, nonatomic measure space and let $(X, |.|_X)$ be a Banach space. We denote by $L^1(P, X)$ the Banach space of all (equivalence classes of) Bochner integrable functions $u: P \to X$ endowed with the norm

$$|u|_{L^1(P,X)} = \int_P |u(t)|_X d\mu$$

A nonempty set $K \subset L^1(P, X)$ is called decomposable if, for every $u, v \in K$ and every $A \in \mathcal{F}$, one has

$$\chi_A.u + \chi_{P \setminus A}.v \in K$$

where $\chi_B, B \in \mathcal{F}$ indicates the characteristic function of B.

A metric space Z is called an absolute retract if, for any metric space X and any nonempty closed set $X_0 \subset X$, every continuous function $g: X_0 \to Z$ has a continuous extension $g: X \to Z$ over X. It is obvious that every continuous image of an absolute retract is an arcwise connected space.

In what follows we recall some preliminary results that are the main tools in the proof of our result.

Let (P, \mathcal{F}, μ) be a finite, positive, nonatomic measure space, S a separable Banach space and let $(X, |.|_X)$ be a real Banach space. To simplify the notation we write E in place of $L^1(P, X)$.

Lemma 2.1 ([12]) Assume that $\phi: S \times E \to 2^E$ and $\psi: S \times E \times E \to 2^E$ are Hausdorff continuous multifunctions with nonempty, closed, decomposable values, satisfying the following conditions

a) There exists $L \in [0, 1)$ such that, for every $s \in S$ and every $u, u' \in E$,

$$D_E(\phi(s, u), \phi(s, u')) \le L|u - u'|_E$$

b) There exists $M \in [0,1)$ such that L + M < 1 and for every $s \in S$ and every $(u, v), (u', v') \in E \times E$,

$$D_E(\psi(s, u, v), \psi(s, u', v')) \le M(|u - u'|_E + |v - v'|_E).$$

Set $Fix(\Gamma(s,.)) = \{u \in E; u \in \Gamma(s,u)\}$, where $\Gamma(s,u) = \psi(s,u,\phi(s,u))$, $(s,u) \in S \times E$. Then

1) For every $s \in S$ the set $Fix(\Gamma(s, .))$ is nonempty and arcwise connected.

2) For any $s_i \in S$, and any $u_i \in Fix(\Gamma(s,.)), i = 1, ..., p$ there exists a continuous function $\gamma : S \to E$ such that $\gamma(s) \in Fix(\Gamma(s,.))$ for all $s \in S$ and $\gamma(s_i) = u_i, i = 1, ..., p$.

Lemma 2.2 ([13]) Let $U: P \to 2^X$ and $V: P \times X \to 2^X$ be two nonempty closed-valued multifunctions satisfying the following conditions

a) U is measurable and there exists $r \in L^1(P)$ such that $D_X(U(t), \{0\}) \leq r(t)$ for almost all $t \in P$.

b) The multifunction $t \to V(t, x)$ is measurable for every $x \in X$.

c) The multifunction $x \to V(t, x)$ is Hausdorff continuous for all $t \in P$.

Let $v: P \to X$ be a measurable selection from $t \to V(t, U(t))$.

Then there exists a selection $u \in L^1(P, X)$ such that $v(t) \in V(t, u(t)), t \in P$.

In the sequel J = [0, T], T > 0 and $\{A(t); t \in J\}$ is the infinitesimal generator of the strongly continuous evolution system G(t, s), $0 \le s \le t \le T$.

Recall that a family of bounded linear operators G(t, s) on X, $0 \le s \le t \le T$ depending on two parameters is said to be a strongly continuous evolution system if there are fulfilled the following conditions: G(s, s) = I, G(t, r)G(r, s) = G(t, s)for $0 \le s \le r \le t \le T$ and $(t, s) \to G(t, s)$ is strongly continuous for $0 \le s \le t \le T$, i.e., $\lim_{t\to s, t>s} G(t, s)x = x$ for all $x \in X$.

Consider the evolution inclusion

$$x'(t) \in A(t)x(t) + \int_0^t K(t,s)F(s,x(s))ds, \quad x(0) = x_0,$$
(2.1)

where $F: J \times X \to \mathcal{P}(X)$ is a set-valued map, X is a separable Banach space, A(t) is the infinitesimal generator of a strongly continuous evolution system of a two parameter family $\{G(t,\tau), t \geq 0, \tau \geq 0\}$ of bounded linear operators of X into $X, D = \{(t, s) \in J \times J; t \ge s\}, K(., .) : D \to \mathbf{R}$ is continuous and $x_0 \in X$. A continuous mapping $x(.) \in C(J, X)$ is called a *mild solution* of problem

(2.1) if there exists a (Bochner) integrable function $f(.) \in L^1(J, X)$ such that

$$f(t) \in F(t, x(t)) \quad a.e. (J) \tag{2.2}$$

$$x(t) = G(t,0)x_0 + \int_0^t G(t,\tau) \int_0^\tau K(\tau,s)f(s)dsd\tau \quad \forall t \in J.$$
 (2.3)

In this case we shall call (x(.), f(.)) a trajectory-selection pair of (2.1). We note that condition (2.3) can be rewritten as

$$x(t) = G(t,0)x_0 + \int_0^t U(t,s)f(s)ds \quad \forall t \in J,$$
(2.4)

where $U(t,s) = \int_s^t G(t,\tau) K(\tau,s) d\tau$. Denote $M := \sup_{t,s \in J} |G(t,s)|$ and $M_0 := \sup_{(t,s) \in D} |K(t,s)|$ and remark that $|U(t,s)| \le MM_0(t-s) \le MM_0T.$

We shall use the following notations for the solution sets of problem (1.1).

 $\mathcal{S}(x_0) = \{x(.); x(.) \text{ is a mild solution of } (1.1)\}.$ (2.5)

In order to study problem (1.1) we introduce the following hypothesis.

Hypothesis 2.3 $F: J \times X \times X \to \mathcal{P}(X)$ and $H: J \times X \to \mathcal{P}(X)$ are two set-valued maps with nonempty closed values, satisfying

i) The set-valued maps $t \to F(t, u, v)$ and $t \to H(t, u)$ are measurable for all $u, v \in X$.

ii) There exist $l(.) \in L^1(J, R)$ such that, for every $u, u' \in X$,

$$D(H(t, u), H(t, u')) \le l(t)|u - u'|$$
 a.e. (J).

iii) There exist $m(.) \in L^1(J, R)$ and $\theta \in [0, 1)$ such that, for every $u, v, u', v' \in$ Χ,

$$D(F(t, u, v), F(t, u', v')) \le m(t)|u - u'| + \theta|v - v'| \quad a.e. (J).$$

iv) There exist $f, g \in L^1(J, R)$ such that

$$d(0,F(t,0,0)) \leq f(t), \quad d(0,H(t,0)) \leq g(t) \quad a.e. \ (J).$$

In what follows $N(t) = \max\{l(t), m(t)\}, t \in J$ and $N^*(t) = \int_0^t N(s) ds$.

Given $\alpha \in R$ we denote by L^1 the Banach space of all (equivalence classes of) Lebesgue measurable functions $\sigma: J \to X$ endowed with the norm

$$|\sigma|_1 = \int_0^T e^{-\alpha N^*(t)} |\sigma(t)| dt.$$

3 The main result

Even if the multifunction from the right-hand side of (1.1) has, in general, nonclosed nonconvex values, the solution set $S(x_0)$ defined in (2.5) has some meaningful properties, stated in theorems below.

Theorem 3.1 Consider $\{A(t); t \in J\}$ the infinitesimal generator of the strongly continuous evolution system $G(t, s), 0 \le s \le t \le T$, assume that F and H satisfy Hypothesis 2.3 and let $\alpha > \frac{2MM_0T}{1-\theta}$. Then 1) For every $x_0 \in X$, the solution set $S(x_0)$ of (1.1) is nonempty and arcwise

1) For every $x_0 \in X$, the solution set $S(x_0)$ of (1.1) is nonempty and arcwise connected in the space C(J,X).

2) For any $\xi_i \in X$ and any $x_i \in S(\xi_i)$, i = 1, ..., p, there exists a continuous function $s : X \to C(J, X)$ such that $s(\xi) \in S(\xi)$ for any $\xi \in X$ and $s(\xi_i) = x_i, i = 1, ..., p$.

3) The set $S = \bigcup_{\xi \in X} S(\xi)$ is arcwise connected in C(J,X).

Proof. 1) For $\xi \in X$ and $v \in L^1$, set

$$x_{\xi}(t) = G(t)\xi + \int_{0}^{t} U(t,s)v(s)ds, \quad t \in J,$$
(3.1)

where $U(t,s) = \int_s^t G(t,\tau) K(\tau,s) d\tau$ and consider $P: X \to C(J,X)$ defined by $P(\xi)(t) = G(t,0)\xi$.

We prove that the multifunctions $\phi: X \times L^1 \to 2^{L^1}$ and $\psi: X \times L^1 \times L^1 \to 2^{L^1}$ given by

$$\phi(\xi, u) = \{ v \in L^1; \quad v(t) \in H(t, x_{\xi}(t)) \quad a.e.(J) \},$$

$$\psi(\xi, u, v) = \{ w \in L^1; \quad w(t) \in F(t, x_{\xi}(t), v(t)) \quad a.e.(J) \},$$

 $\varphi(\mathbf{s}, \omega, v) \quad (\omega = 2, \omega (v) = -(v, \omega \zeta(v), v(v)) \quad \omega$

 $\xi \in X, \, u,v \in L^1$ satisfy the hypotheses of Lemma 2.1.

Since $x_{\xi}(.)$ is measurable and H satisfies Hypothesis 2.3 i) and ii), the multifunction $t \to H(t, x_{\xi}(t))$ is measurable and nonempty closed-valued, it has a measurable selection. Therefore due to Hypothesis 2.3 iv), the set $\phi(\xi, u)$ is nonempty. The fact that the set $\phi(\xi, u)$ is closed and decomposable follows by a simple computation. In the same way we obtain that $\psi(\xi, u, v)$ is a nonempty closed decomposable set.

Pick $(\xi, u), (\xi_1, u_1) \in X \times L^1$ and choose $v \in \phi(\xi, u)$. For each $\varepsilon > 0$ there exists $v_1 \in \phi(\xi_1, u_1)$ such that, for every $t \in J$, one has

$$|v(t) - v_1(t)| \le D(H(t, x_{\xi}(t)), H(t, x_{\xi_1}(t))) + \varepsilon \le l(t)[M|\xi - \xi_1| + MM_0T \int_0^t |u(s) - u_1(s)|ds] + \varepsilon.$$

Hence

$$|v - v_1|_1 \le M |\xi - \xi_1| \int_0^T e^{-\alpha N^*(t)} l(t) dt + M M_0 T \int_0^T e^{-\alpha N^*(t)} l(t) (\int_0^t |u(s)| dt + M M_0 T \int_0^T e^{-\alpha N^*(t)} l(t) (\int_0^t |u(s)| dt + M M_0 T \int_0^T e^{-\alpha N^*(t)} l(t) dt + M M_0 T \int_0^T e^{-\alpha N^*(t)} l(t) (\int_0^t |u(s)| dt + M M_0 T \int_0^T e^{-\alpha N^*(t)} l(t) dt + M M_0 T$$

$$-u_1(s)|ds)dt + \varepsilon T \le \frac{M}{\alpha}|\xi - \xi_1| + \frac{MM_0T}{\alpha}|u - u_1|_1 + \varepsilon T$$

for any $\varepsilon > 0$.

This implies

$$d_{L^1}(v,\phi(\xi_1,u_1)) \le \frac{M}{\alpha} |\xi - \xi_1| + \frac{MM_0T}{\alpha} |u - u_1|_1$$

for all $v \in \phi(\xi, u)$. Therefore,

$$d_{L^1}^*(\phi(\xi, u), \phi(\xi_1, u_1)) \le \frac{M}{\alpha} |\xi - \xi_1| + \frac{MM_0T}{\alpha} |u - u_1|_1.$$

Consequently,

$$D_{L^1}(\phi(\xi, u), \phi(\xi_1, u_1)) \le \frac{M}{\alpha} |\xi - \xi_1| + \frac{MM_0T}{\alpha} |u - u_1|_1.$$

which shows that ϕ is Hausdorff continuous and satisfies the assumptions of Lemma 2.1.

Pick $(\xi, u, v), (\xi_1, u_1, v_1) \in X \times L^1 \times L^1$ and choose $w \in \psi(\xi, u, v)$. Then, as before, for each $\varepsilon > 0$ there exists $w_1 \in \psi(\xi_1, u_1, v_1)$ such that for every $t \in J$

$$|w(t) - w_1(t)| \le D(F(t, x_{\xi}(t), v(t)), F(t, x_{\xi_1}(t), v_1(t))) + \varepsilon \le$$
$$m(t)[M|\xi - \xi_1| + MM_OT \int_0^t |u(s) - u_1(s)|ds] + \theta |v(t) - v_1(t)| + \varepsilon.$$

Hence

$$\begin{split} |w - w_1|_1 &\leq \frac{M}{\alpha} |\xi - \xi_1| + \frac{MM_0T}{\alpha} |u - u_1|_1 + \theta |v - v_1|_1 + \varepsilon T \\ &\leq \frac{M}{\alpha} |\xi - \xi_1| + (\frac{MM_0T}{\alpha} + \theta)(|u - u_1|_1 + |v - v_1|_1) + \varepsilon T \\ &\leq \frac{M}{\alpha} |\xi - \xi_1| + (\frac{MM_0T}{\alpha} + \theta)d_{L^1 \times L^1}((u, v), (u_1, v_1)) + \varepsilon T. \end{split}$$

As above, we deduce that

$$\begin{aligned} D_{L^1}(\psi(\xi, u, v), \psi(\xi_1, u_1, v_1)) &\leq \\ \frac{M}{\alpha} |\xi - \xi_1| + (\frac{MM_0T}{\alpha} + \theta) d_{L^1 \times L^1}((u, v), (u_1, v_1)). \end{aligned}$$

namely, the multifunction ψ is Hausdorff continuous and satisfies the hypothesis of Lemma 2.1.

Define $\Gamma(\xi, u) = \psi(\xi, u, \phi(\xi, u)), (\xi, u) \in X \times L^1$. According to Lemma 2.1, the set $Fix(\Gamma(\xi, .)) = \{u \in L^1; u \in \Gamma(\xi, u)\}$ is nonempty and arcwise connected in $L^1(J, X)$. Moreover, for fixed $\xi_i \in X$ and $u_i \in Fix(\Gamma(\xi_i, .)), i = 1, ..., p$, there exists a continuous function $\gamma: X \to L^1$ such that

$$\gamma(\xi) \in Fix(\Gamma(\xi, .)), \quad \forall \xi \in X, \tag{3.2}$$

114

Mild solutions of a evolution inclusion

$$\gamma(\xi_i) = u_i, \quad i = 1, ..., p.$$
 (3.3)

We shall prove that

$$Fix(\Gamma(\xi,.)) = \{ u \in L^1; \quad u(t) \in F(t, x_{\xi}(t), H(t, x_{\xi}(t))) \quad a.e. \ (J) \}.$$
(3.4)

Denote by $A(\xi)$ the right-hand side of (3.4). If $u \in Fix(\Gamma(\xi, .))$ then there is $v \in \phi(\xi, v)$ such that $u \in \psi(\xi, u, v)$. Therefore, $v(t) \in H(t, x_{\xi}(t))$ and

$$u(t) \in F(t, x_{\xi}(t), v(t)) \subset F(t, x_{\xi}(t), H(t, x_{\xi}(t)))$$
 a.e. (J),

so that $Fix(\Gamma(\xi, .)) \subset A(\xi)$.

Let now $u \in A(\xi)$. By Lemma 2.2, there exists a selection $v \in L^1$ of the multifunction $t \to H(t, x_{\xi}(t))$ satisfying

$$u(t) \in F(t, x_{\xi}(t), v(t))$$
 a.e. (J)

Hence, $v \in \phi(\xi, v)$, $u \in \psi(\xi, u, v)$ and thus $u \in \Gamma(\xi, u)$, which completes the proof of (3.4).

We next note that the function $W: L^1 \to C(J, X)$,

$$W(v)(t):=\int_0^t U(t,s)v(s)ds$$

is continuous and one has

$$\mathcal{S}(\xi) = P(\xi) + W(Fix(\Gamma(\xi, .))), \quad \xi \in X.$$
(3.5)

Since $Fix(\Gamma(\xi, .))$ is nonempty and arcwise connected in $L^1(J, X)$, the set $S(\xi)$ has the same properties in C(J, X).

2) Let $\xi_i \in X$ and let $x_i \in \mathcal{S}(\xi_i), i = 1, ..., p$ be fixed. By (3.5) there exists $v_i \in Fix(\Gamma(\xi_i, .))$ such that

$$x_i = P(\xi_i) + W(v_i), \quad i = 1, ..., p.$$

If $\gamma : X \to L^1$ is a continuous function satisfying (3.2) and (3.3) we define, for every $\xi \in X$,

$$s(\xi) = P(\xi) + W(\gamma(\xi)).$$

Obviously, the function $s: X \to C(J, X)$ is continuous, $s(\xi) \in \mathcal{S}(\xi)$ for all $\xi \in X$ and

$$s(\xi_i) = P(\xi_i) + W(\gamma(\xi_i)) = P(\xi_i) + W(v_i) = x_i, \quad i = 1, ..., p.$$

3) Let $x_1, x_2 \in \mathcal{S} = \bigcup_{\xi \in X} \mathcal{S}(\xi)$ and choose $\xi_i \in X$, i = 1, 2 such that $x_i \in S(\xi_i)$, i = 1, 2. From the conclusion of 2) we deduce the existence of a continuous function $s : X \to C(J, X)$ satisfying $s(\xi_i) = x_i$, i = 1, 2 and $s(\xi) \in \mathcal{S}(\xi)$, $\xi \in X$. Let $h : [0, 1] \to X$ be a continuous mapping such that $h(0) = \xi_1$ and $h(1) = \xi_2$. Then the function $s \circ h : [0, 1] \to C(J, X)$ is continuous and verifies

$$s \circ h(0) = x_1, \quad s \circ h(1) = x_2, \quad s \circ h(\tau) \in \mathcal{S}(h(\tau)) \subset \mathcal{S}, \quad \tau \in [0, 1].$$

Remark 3.2 If $A(t) \equiv A$ and A is the infinitesimal generator of a strongly continuous semigroup of bounded linear operators $\{G(t); t \geq 0\}$ from X to X then problem (1.1) reduces to the problem

$$x'(t) \in Ax(t) + \int_0^t K(t,s)F(s,x(s),H(s,x(s)))ds, \quad x(0) = x_0.$$
(3.6)

Qualitative properties of the solution set of problem (3.6) may be found in [1-7,11] etc.

Obviously, a similar result to the one in Theorem 3.1 may be obtained for problem (3.6).

References

- A. ANGURAJ, C. MURUGESAN, Continuous selections of set of mild solutions of evolution inclusions, *Electronic J. Diff. Equations*, **2005** (2005), No. 21, 1-7.
- [2] K. BALACHANDRAN, P. BALASUBRAMANIAN, J.P. DAUER, Controllability of nonlinear integrodifferential systems in Banach spaces, J. Optim. Theory Appl., 74 (1995), 83-91.
- [3] M. BENCHOHRA, S. K. NTOUYAS, Existence results for neutral functional differential and integrodifferential inclusions in Banach spaces, *Electronic J. Diff. Equations*, **2000** (2000), No. 20, 1-15.
- [4] M. BENCHOHRA, S. K. NTOUYAS, Nonlocal Cauchy problems for neutral functional differential and integrodifferential inclusions in Banach spaces, J. Math. Anal. Appl., 258 (2001), 573-590.
- [5] M. BENCHOHRA, S. K. NTOUYAS, Controllability of infinite time horizon for functional differential and integrodifferential inclusions in Banach spaces, *Commun. Applied Nonlin. Anal.*, 8 (2001), 63-78.
- [6] M. BENCHOHRA, S. K. NTOUYAS, Existence results for functional differential and integrodifferential inclusions in Banach spaces, *Indian J. Pure Applied Math.*, **32** (2001), 665-675.
- [7] M. BENCHOHRA, S. K. NTOUYAS, Controllability for functional and integrodifferential inclusions in Banach spaces, J. Optim. Theory Appl., 113 (2002), 449-472.
- [8] A. CERNEA, On the set of solutions of some nonconvex nonclosed hyperbolic differential inclusions, *Czechoslovak Math. J.*, **52(127)** (2002), 215-224.
- [9] A. CERNEA, On the solution set of a nonconvex nonclosed higher order differential inclusion, *Math. Commun.*, **12** (2007), 221-228.

116

- [10] A. CERNEA, On the solution set of some classes of nonconvex nonclosed differential inclusions, *Portugaliae Math.*, to appear.
- [11] B. LIU, Controllability of neutral functional differential and integrodifferential inclusions with infinite delay, J. Optim. Theory Appl., 123 (2004), 573-593.
- [12] S. MARANO, Fixed points of multivalued contractions with nonclosed, nonconvex values, Atti. Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur., 5 (1994), 203-212.
- [13] S. MARANO, V. STAICU, On the set of solutions to a class of nonconvex nonclosed differential inclusions, *Acta Math. Hungar.*, **76** (1997), 287-301.

Received: 04.10.2008, Revised: 23.10.2008, Accepted: 11.05.2010.

> Faculty of Mathematics and Informatics, University of Bucharest, Academiei 14, 010014 Bucharest, Romania. E-mail: acernea@fmi.unibuc.ro